

**University Code: 10384**

**Student ID No.: 20620071154072**

廈門大學

**MASTER DEGREE THESIS**

**Biosynthesis of Silver Nanoparticles by sundried *Aleurites moluccana* leaf and yeast *pichia pastoris***

Supported by the National Natural Science Foundation of China (Grant No. 20776120 and 20576109), the National High Technology Research and Development Program of China (863 Program, Grant No. 2007AA03Z347) and the Natural Science Foundation of Fujian Province of China (Grant No. 2008J0169)

By  
**Egesa Dan**  
(Uganda)

**Supervisor: Professor Li Qingbiao**

**Program: Chemical Engineering**

**Date of Graduation: July, 2009**

A THESIS SUBMITTED IN FULFILLMENT OF THE REQUIREMENTS FOR THE  
AWARD OF THE DEGREE OF MASTER OF ENGINEERING  
DEPARTMENT OF CHEMICAL AND BIOCHEMICAL ENGINEERING  
COLLEGE OF CHEMISTRY AND CHEMICAL ENGINEERING  
XIAMEN UNIVERSITY

厦门大学博硕士论文摘要库

# CERTIFICATION

I, Professor \_\_\_\_\_, hereby certify that I have read this manuscript and recommend for acceptance by the Xiamen University a dissertation entitled “Biosynthesis of silver nanoparticles by sundried *Aleurites moluccana* leaf and yeast *pichia pastoris*” in fulfillment of degree of Master of Engineering at Xiamen University, People’s Republic of China.

Signed.....

Supervisor

Date.....

Department of Chemical and Biochemical Engineering

College of Chemistry and Chemical Engineering

Xiamen University

Xiamen, Fujian Province

P.R. China

## **ORIGINAL STATEMENT**

The research described in this Master of Engineering thesis was conducted under the supervision of Professor Professor Li Qingbiao at the Department of Chemical and Biochemical Engineering, Xiamen University. I hereby declare that the work submitted is my own and that appropriate credit has been given where reference has been made to the work of others. I also confirm that it has not been previously or concurrently submitted for any other degree, diploma or any other qualifications at Xiamen University, P.R China or other institutions.

## **COPYRIGHT DECLARATION**

All rights reserved. No part of this dissertation may be reproduced, stored in any retrieval system, or transmitted in any form by any means: electronic, mechanical, photocopying, recording or otherwise without prior written permission of the author supervisor and or Xiamen University.

---

EGESA DAN

Date:

## **ACKNOWLEDGEMENT**

All thanks and praise to God Almighty for the gift of life and grace to do this work without which it wouldn't be possible. This work is dedicated to Him.

I would like to thank the National Natural Science Foundation of China (Grant No. 20776120 and 20576109), the National High Technology Research and Development Program of China (863 Program, Grant No. 2007AA03Z347) and the Natural Science Foundation of Fujian Province of China (Grant No. 2008J0169) that supported this research.

I would like to thank everyone who assisted me over the course of my research project. In particular I would like to thank my supervisor professor Li Qingbiao for his dedicated and wise advice.

I am also grateful to all of the students who helped in peer review specifically Li Qin, Jiale Huang and Zhou Yao.

I would also like to thank my parents Captain John Egesa and Harriet Egesa for there support and my wife Ritah Egesa for he encouragement.

## ABSTRACT

Because the properties of silver nanoparticles depend so closely on their size, size distribution and morphology, controlling the size and shape of silver nanoparticles is of great interest to researchers today. In this study, we focus on the control of size of silver nanoparticles biologically synthesized using *aleurites moluccana* leaf broth and *pichia pastoris*, as reducing agents. The effect of reaction temperature, rotate rate, the ratio between biomass concentration to silver ions, and pH on the synthesis process of silver nanoparticles was investigated. The formation of nanoparticles was understood and monitored by UV-Vis spectrometer, transmission electron microscopy (TEM), scanning electron microscopy (SEM), and Fourier transform infrared spectrometer (FTIR).

Using *A.moluccana* leaf broth as reducing agent, when the reaction temperature was fixed at 80 °C, the mean particle diameter was 31.1 nm, increase in reaction temperature to 100 °C, led to reduction of the mean particle diameter to 13.2 nm. Increasing the rotate rate led to steady reduction in absorbance and a blue shift in the absorbance peak, there was reduction in particle size from 26.3 nm to 25.5 nm with increase in rotate rate from 10 rpm to 30 rpm respectively. While increasing biomass concentration led to increase in absorbance and a blue shift in the position of the spectrum suggesting reduction in particle size. In addition, pH was found to have a great impact on the size of silver nanoparticle. When pH of the biomass solution was increased from 5 to 9 to 12, this led to a reduction in particle size from 11.9 nm to 11.3 nm and to 8.7 nm respectively.

Using *P.pastoris* biomass as reducing agent, silver nitrate and silver amine were used in separate experiments as sources of silver ions. A comparison was made between these two sources of silver ions under similar reaction conditions, when silver amine was used the bioreduction was much faster and the absorbance peak was higher compared to silver nitrate. When silver amine was source of silver ions, in acidic condition at pH 4 and below there was no bioreduction and increase in pH led to reduction in particle size. When silver nitrate was used as source of silver ions, there was also no bioreduction in acidic conditions at pH 4 and below. Increase in pH led to steady increase in absorbance and a blue shift in the position of absorbance

peak from 423 nm at pH 5 to 418 nm at pH 12. Increasing the rotate rate and temperature led to a steady increase in the absorbance and there was no shift in position of the absorbance peak. Increasing biomass concentration led to increase in particle size from 11.98 nm to 12.97 nm. Increasing the ratio of silver ions to biomass led to increase in absorbance and broadening of peak hence suggesting aggregation of nanoparticles in solution.

**KEY WORDS:** Biosynthesis; Nanoparticles; Silver; *Aleurites moluccana*; *Pichia pastoris*

## 摘要

由于纳米银的性质和它的粒径、粒径分布和形貌息息相关，因此如何调控纳米银的粒径和形貌引起了研究者的广泛关注。本研究选取了两种生物质——石栗叶和毕赤酵母菌粉作为还原剂来制备纳米银，研究了反应时间、温度、搅拌速度、生物质/银离子的比值和 pH 值等条件对纳米银的粒径和形貌的影响，并利用紫外-可见分光光度计 (UV-Vis)、扫描电子显微镜 (SEM)、透射电子显微镜 (TEM) 和傅立叶变换红外光谱仪 (FTIR) 等多种分析方法来表征所形成的纳米银。

利用石栗叶作为还原剂，80 °C 时所制备的纳米银的平均粒径大约为 31.1 nm，当温度升高到 100 °C 时，所合成的纳米银的平均粒径减小为 13.2 nm。增大搅拌速度使得纳米银的紫外-可见吸收峰的位置发生了蓝移，峰的强度减弱，当搅拌速度从 10 rpm 增大到 30 rpm，所获得的纳米银的平均粒径有轻微减小 (26.3 nm→25.5 nm)。而增加石栗叶的浓度则会使得紫外-可见吸收峰的峰强变大，峰的位置发生了蓝移，这说明了随着石栗叶浓度的增加，所制得的纳米银的粒径逐渐减小。此外，我们发现了 pH 值对于纳米银的粒径也有很大的影响，当溶液的 pH 逐渐增大时 (5→9→12)，所制得的纳米银的粒径逐渐减小 (11.9 nm→11.3 nm→8.7 nm)。

利用毕赤酵母菌粉作为还原剂时，本研究选取了硝酸银和银氨来作为银前驱体，比较了二者的差异，研究发现利用银氨作为前驱体时的还原速率要比用硝酸银时快。利用硝酸银作为前驱体时，当  $\text{pH} \leq 4$  时，溶液中没有纳米银生成；当  $\text{pH} > 4$  时，随着 pH 值的增大，所获得的纳米颗粒的粒径逐渐减小。而利用银氨作为前驱体时， $\text{pH} \leq 4$  时，溶液中同样没有纳米银生成，随着 pH 值的增大 (5→12)，纳米银的紫外-可见吸收峰逐渐蓝移 (423 nm→418 nm)，说明了纳米银的粒径逐渐减小。随着搅拌速度的增大和温度的提高，纳米银的紫外-可见吸收峰的强度逐渐增大，但峰的位置没有发生变化，说明搅拌速度和温度对纳米银的粒径影响不大。银离子浓度一定时，所制得的纳米银的平均粒径随着毕赤酵母菌粉量的增大有轻微的增大 (11.98 nm→12.97 nm)。而生物质浓度一定时，增大硝酸银的浓度，所制得的纳米银的紫外-可见吸收峰强度增大，但半峰宽明显变宽，说明



溶液中纳米银发生了团聚。

**关键词：** 生物合成；纳米颗粒；银；石栗；毕赤酵母

厦门大学博硕士论文摘要库

## LIST OF ABBREVIATIONS

fcc	Face Centered Cubic
fig.	Figure
FTIR	Fourier Transform Infrared
g	Gram
h	Hour
M	Molar
min	Minute
L	Microlitre
mM	Millimolar
nm	Nanometre
rpm	Revolution Per Minute
SEM	Scanning Electron Microscopy
TEM	Transmission Electron Microscopy
UV-Vis	Ultraviolet-Visible
XRD	X-ray Diffraction
a.u	Arbitrary Units
abs	Absorbance
gp	Group

# TABLE OF CONTENTS

<b>ACKNOWLEDGEMENT</b> .....	I
<b>ABSTRACT</b> .....	III
<b>摘要</b> .....	V
<b>LIST OF ABBREVIATIONS</b> .....	VII
<b>TABLE OF CONTENTS</b> .....	IX
<b>LIST OF FIGURES</b> .....	XIII
<b>LIST OF TABLES</b> .....	XIX
<b>CHAPTER 1 INTRODUCTION AND LITERATURE SURVEY</b> .....	1
1.1. Introduction and Emergence of Nanotechnology .....	1
1.1.1. What is nanoscience and nanotechnology? .....	1
1.1.2. Emergence of Nanotechnology .....	1
1.2. Properties and Applications of silver nanoparticles .....	2
1.2.1 Properties of silver nanoparticles .....	2
1.2.2. Applications of silver nanoparticles. ....	2
1.2.2.1. Antibacterial applications .....	2
1.2.2.2. Antiviral applications .....	3
1.2.2.3. Medical applications .....	3
1.2.2.4. Catalytic applications .....	3
1.2.2.5. Other applications .....	4
1.3. Preparation methods of silver nanoparticles .....	4
1.3.1. Chemical methods .....	5
1.3.1.1. Reduction of silver ions in the presence of reducing agent $\text{NaBH}_4$ .....	5
1.3.1.2. Reduction of silver nitrate with ethylene glycol .....	6
1.3.1.3. Synthesis of Nanoparticles by Pyrolysis .....	7
1.3.1.4. The reverse micelle method .....	8
1.3.1.5. Sonochemical method .....	8
1.3.2. Physical methods .....	9
1.3.3. Biological methods .....	10
1.3.3.1. Using microorganisms .....	10

1.3.3.2. Using Plant Biomass.....	15
1.4. Factors affecting the biosynthesis process of silver nanoparticles .....	16
1.4.1 Effect of Temperature.....	16
1.4.2 Effect of pH.....	17
1.4.3 Effect of biomass concentration.....	17
1.4.4 Effect of Reaction Time .....	17
1.5. Aim, significance and content of this study.....	18
<b>CHAPTER 2 MATERIALS AND METHODS.....</b>	<b>19</b>
2.1. Reagents and instruments .....	19
2.2 Biosynthesis of silver nanoparticles using <i>Aleurites moluccana</i> leaves.....	19
2.2.1 Preparation of <i>A.moluccana</i> leaf biomass and reagents.....	20
2.2.2. Preparation of silver nanoparticles.....	20
2.2.2.1 Preparation of silver nanoparticles at different pH.....	20
2.2.2.2 Preparation of silver nanoparticles at different <i>A.moluccana</i> biomass concentrations. .....	20
2.2.2.3 Preparation of silver nanoparticles at different temperature.....	21
2.2.2.4 Preparation of silver nanoparticles at different rotate rate.....	21
2.3 Biosynthesis of silver nanoparticles using yeast <i>pichia pastoris</i> GS115.....	21
2.3.1 Preparation of <i>P.pastoris</i> biomass and reagents .....	21
2.3.2 Preparation of silver nanoparticles.....	22
2.3.2.1 Preparation of silver nanoparticles at different pH values.....	22
2.3.2.2 Preparation of silver nanoparticles at different <i>P.pastoris</i> biomass concentration ..	22
2.3.2.3 Preparation of silver nanoparticles at different temperature.....	22
2.3.2.4 Preparation of silver nanoparticles at different rotate rates .....	22
2.4 Methods of characterization.....	23
2.4.1 Visual Inspection .....	23
2.4.2 UV-Vis spectra analysis.....	24
2.4.3 Fourier Transform infrared spectroscopy (FTIR) analysis.....	24
2.4.4 TEM characterization of silver nanoparticles .....	25
2.4.5 SEM characterization .....	25
<b>CHAPTER 3 BIOSYNTHESIS OF SILVER NANOPARTICLES USING <i>ALEURITES</i></b>	

<b>MOLUCCANA LEAF BIOMASS</b> .....	26
3.1. Introduction.....	26
3.2 Characterization of silver nanoparticles synthesized by <i>A.moluccana</i> leaf.....	27
3.2.1. UV-Vis spectroscopy results .....	27
3.2.2. TEM results .....	28
3.2.3. FTIR spectroscopy results .....	29
3.3 Effect of synthesizing conditions on the synthesis process of silver nanoparticles.....	31
3.3.1. Effect of reaction time.....	31
3.3.2. Effect of pH on the synthesis process of silver nanoparticles.....	36
3.3.3. Effect of biomass concentration .....	42
3.3.3.1. Volume of 0.1M silver nitrate fixed at 1ml.....	42
3.3.3.2. Volume of 0.1 M silver nitrate fixed at 2 ml.....	45
3.3.3.3. Volume of 0.1 M silver nitrate fixed at 3 ml.....	48
3.3.3.4. Volume of 0.1 M silver nitrate fixed at 4 ml.....	51
3.3.4. Effect of temperature.....	53
3.3.5. Effect of rotate rate.....	57
.....	59
3.4. Summery .....	64
<b>CHAPTER 4 BIOSYNTHESIS OF SILVER NANOPARTICLES USING DEAD YEAST</b>	
<b>(<i>Pichia Pastoris</i>)</b> .....	65
4.1 Introduction.....	65
4.2. Characterization of silver nanoparticles synthesized by <i>pichia pastoris</i> .....	65
4.2.1. UV-VIS spectroscopy results .....	65
4.2.3. Fourier transform infrared spectroscopy (FTIR).....	67
4.2.3.1 FTIR analysis of dried biomass of <i>pichia pastoris</i> before and after bioreduction ..	67
4.2.3.2. FTIR analysis of silver nanoparticles produced by <i>P.pastoris</i> biomass .....	68
4.3. Effect of synthesizing conditions on the synthesis process of silver Nanoparticles.....	69
4.3.1. Effect of reaction time.....	69
4.3.2. Effect of pH.....	71
4.3.2.1. Using 0.1 M silver nitrate solution as source of silver ions.....	71
4.3.2.2. Using 0.1 M silver Amine solution as source of silver ions. ....	75

4.3.3. Effect of biomass concentration .....	78
4.3.4. Effect of temperature.....	83
4.3.5. Effect of rotate rate.....	85
4.3.6. Effect of silver amine ratio to biomass.....	87
4.4. Summary .....	90
<b>CONCLUSION AND SUGGESTIONS .....</b>	<b>91</b>
<b>References.....</b>	<b>93</b>

## LIST OF FIGURES

<b>Figure.1.1</b> (A) Low and (B) high-magnification SEM images of slightly truncated silver nanocubes synthesized with the polyol process.....	6
<b>Figure.1.2.</b> TEM images of silver nanocubes synthesized under different conditions.....	7
<b>Figure 1.3.</b> Flow chart for synthesis of silver nanoparticles by reverse micelle.....	8
<b>Figure.1.4</b> Self-assembled silver nanochains.....	9
<b>Figure1.5</b> , Transmission electron micrograph of a <i>Pseudomonas stutzeri</i> AG259 cell grown on a 50 mM Ag <sup>+</sup> -containing agar substrate. Silver-based single crystals with different size and morphology are associated with the cell. ....	12
<b>Figure.1.6</b> above A variety of silver (Ag) crystal morphologies. The particles are a selection of different crystal morphologies that are found in whole cell preparations and thin sections. ....	12
<b>Figure.3.1</b> shows UV-Vis spectra recorded with respect to time after the reaction of 3 ml of 0.1 M AgNO <sub>3</sub> solution with 1 g of <i>aleurites moluccana</i> powder in 100ml of distilled water at temperature of 80 °C and rotate rate of 30 rpm.....	27
<b>Figure.3.2</b> , representative TEM pictures recorded from the silver nanoparticle. The nanoparticles were produced using 1 g of <i>aleurites moluccana</i> leaf biomass as reducing agent, 3 ml of 0.1 M silver nitrate solution as source of silver ions. ....	28
<b>Figure.3.3</b> Histogram showing the size distribution of silver nanoparticles synthesized using 1 g of <i>A.moluccana</i> . Most of the nanoparticles are in the size range of between 30-60 nm. The average particle diameter is 48.8 nm.....	29
<b>Figure 3.4</b> FTIR Spectra.....	30
<b>Figure.3.5</b> shows UV-Vis spectra recorded with respect to time after the reaction of 3 ml of 0.1 M AgNO <sub>3</sub> solution with 1 g of <i>aleurites moluccana</i> powder in 100 ml of distilled water at a temperature of 80°C and rotate rate of 30 rpm.....	31
<b>Figure 3.6</b> showing the effect of reaction time (a) on the absorbance peak, (b) on the conversion of silver nanoparticles produced by 1 g of <i>A.moluccana</i> leaf biomass. ....	32
<b>Figure 3.7</b> (a) shows the effect of reaction time on the wavelength, (b), on the size of silver nanoparticles produced by 1 g of <i>A.moluccana</i> biomass. ....	33
<b>Figures.3.8</b> , (a), (b) and (c) show SEM images of silver nanoparticles synthesized.....	34

<b>Figure.3.9</b> Histogram showing particle size distribution of silver nanoparticles synthesized using 1g of <i>A.moluccana</i> biomass and 3 ml of 0.1M silver nitrate at a temperature of 80 °c and a rotate rate of 30 rpm, (a) after 30 minutes of reaction, (b) after 90 minutes of reaction, and (c) after 150 minutes. ....	35
<b>Figure 3.10.</b> Shows UV-Vis spectrum of silver nanoparticles with time at different pH values....	37
<b>Figure.3.11.</b> UV-Vis spectra showing the effect of pH on size and shape of silver nanoparticles. ....	38
<b>Figure 3.12</b> (a) Shows the trend of pH with absorbance maxima, (b) with number of nanoparticles of silver synthesized using <i>aleurites moluccana</i> biomass solution. ....	39
<b>Figure 3.13.</b> (a) Shows the effect of increase in pH on the wavelength, (b) shows the effect of increase in pH on the particle size of silver nanoparticles. ....	40
<b>Figure 3.14.</b> The images of silver nanoparticles (a) at pH of 12, (b) at pH of 9 and (c) pH.5. ....	41
<b>Figure 3.15.</b> The size distribution of silver nanoparticles (a) at pH.12, (b) at pH 9 and (c) at pH 5. ....	41
<b>Figure 3.16.</b> Absorption spectra of silver nanoparticles at temperature of 80 0c.....	43
<b>Figure.3.17.</b> showing UV-Vis spectrum of silver nanoparticles synthesized by 1 g,2 g,3 g and 4 g of <i>A.moluccana</i> leaf biomass after 2 hours of reaction. ....	44
<b>Figure 3.18</b> (a). Shows the variation of biomass concentration with wavelength, (b) the variation of biomass concentration with absorbance.....	45
<b>Figure 3.19</b> showing the effect of biomass concentration at a fixed volume of silver nitrate of 2 ml and rotate rate of 30 rpm, (a) biomass was fixed at 1 g, (b) biomass was fixed at 2 g, (c) biomass was fixed at 3 g, and (d) biomass was fixed at 4 g.....	46
<b>Figure 3.20</b> , the volume of silver nitrate was kept at 2 ml and biomass concentration was adjusted in separate experiments from 1 g, 2 g, 3 g and 4 g. ....	47
<b>Figure 3.21</b> (a) the trend of absorbance with increase in biomass concentration at 2 ml of 0.1 M silver nitrate solution; (b) illustrates the trend of wavelength with increase in biomass concentration. ....	48
<b>Figure 3.22</b> ,Shows the effect of biomass concentration when the volume of silver nitrate is fixed at 3 ml (a) biomass was fixed at 1 g, (b), biomass was fixed at 2 g, (c), biomass was fixed at 3 g, and (d), biomass was fixed at 4 g. ....	49
<b>Figure 3.23</b> , the volume of 0.1 M silver nitrate was fixed at 3mls, rotate rate at 30 rpm,	



Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to [etd@xmu.edu.cn](mailto:etd@xmu.edu.cn) for delivery details.

厦门大学博硕士论文摘要库